

Communicable Disease Prevention During Emergencies and Disasters






Communicable Disease Prevention During Emergencies and Disasters

Emergencies and disasters may increase the risk of communicable disease outbreaks due to overcrowding, disruption of water and sanitation services, limited access to healthcare, interruption of essential services, population displacement, or living in shelters and temporary accommodation centers. Natural disasters, health emergencies, conflicts, and service interruptions may contribute to the spread of respiratory infections, gastrointestinal illnesses, skin infections, and other communicable diseases, especially among vulnerable groups such as children, older adults, and individuals with chronic illnesses or weakened immune systems.

In addition, stress, fatigue, poor nutrition, and difficult living conditions during emergencies may weaken the body's immune system, increasing the risk of infection and disease transmission within communities. Following preventive measures and official public health guidance can help reduce disease spread and protect individuals and communities during emergencies.



The Importance of Communicable Disease Prevention During Emergencies

During emergencies, crowded shelters, evacuation centers, and gathering areas may become high-risk environments for the spread of infectious diseases due to close contact between individuals and difficulties maintaining proper hygiene and ventilation. Disruptions to water, electricity, sanitation, and waste management services may further increase health risks.

Communicable diseases can spread rapidly in crowded settings, especially when personal items are shared, hand hygiene is poor, or cleaning and disinfection measures are inadequate. Disease outbreaks may place additional pressure on healthcare facilities and disrupt essential services if not controlled early.

Infectious diseases may spread through:

- Respiratory droplets from coughing or sneezing
- Direct contact with infected individuals
- Contaminated food or water
- Shared surfaces and equipment
- Poor personal hygiene and inadequate handwashing

Diseases that may increase during emergencies include:

- Influenza and respiratory infections
- COVID-19 and similar viral illnesses
- Gastrointestinal infections and diarrhea
- Skin infections
- Foodborne and waterborne diseases



Prevention of Respiratory Infections

Respiratory infections spread easily during emergencies, particularly in crowded indoor environments such as shelters, temporary housing facilities, and public transportation. Viruses and bacteria may spread through airborne droplets or by touching contaminated surfaces and then touching the face, nose, or mouth.

Poor ventilation and overcrowding can increase the speed of disease transmission, especially among people with weakened immune systems or chronic health conditions. Some respiratory infections may be mild, while others can cause severe complications in vulnerable individuals.

Symptoms of respiratory infections may include:

- Fever
- Cough
- Sore throat
- Runny nose
- Shortness of breath
- Fatigue and body weakness

Prevention measures:

- Maintain good ventilation whenever possible
- Avoid unnecessary crowding
- Wear masks when symptomatic or in crowded places if recommended by health authorities
- Stay home or remain isolated when feeling unwell
- Avoid sharing personal items

Health-related action to take:

- Monitor respiratory symptoms among family members
- Encourage rest and adequate fluid intake for mild illness
- Use prescribed medications according to medical advice

👉 **Following preventive measures helps reduce transmission within the community.**



Prevention of Foodborne and Waterborne Diseases

Food and water may become contaminated during disasters due to flooding, power outages, sanitation failures, or improper food storage. High temperatures and refrigeration failures may also increase the risk of food spoilage and food poisoning.

Foodborne and waterborne illnesses are among the most common health problems during emergencies, particularly in disaster-affected areas or locations with disrupted services. Infections may spread quickly among family members or within shelters and temporary housing facilities.

Contaminated food or water may cause:

- Diarrhea
- Vomiting
- Food poisoning
- Dehydration
- Gastrointestinal infections

Prevention measures:

- Use safe and approved drinking water
- Wash hands before preparing or eating food
- Store food safely and at proper temperatures
- Discard spoiled or contaminated food
- Wash fruits and vegetables thoroughly

Health-related actions to take:

- Replace fluids lost through vomiting or diarrhea
- Monitor children and older adults for signs of dehydration
- Seek medical care if symptoms persist, fever develops, or blood appears in stool
- Use oral rehydration solutions when needed
- Avoid preparing food while ill

📌 Safe food and water are essential for protecting health during emergencies.

Personal Hygiene and Environmental Cleanliness

Good personal hygiene and environmental cleanliness help reduce the spread of germs and viruses within homes, shelters, and temporary accommodation centers. Maintaining clean surroundings also helps reduce insects, unpleasant odors, and other health hazards that may emerge during crises.

Accumulation of waste, poor ventilation, and inadequate cleaning may increase the risk of communicable diseases, skin infections, and respiratory illnesses, especially in crowded environments.

Important hygiene practices include:

- Frequent handwashing
- Regular bathing and changing clothes
- Cleaning shared surfaces and equipment
- Proper waste disposal
- Ventilating indoor spaces whenever possible

Health-related actions to take:

- Clean and cover wounds properly
- Use separate personal items for sick individuals
- Disinfect frequently touched surfaces
- Wash contaminated clothing with soap and water
- Seek medical attention for infected wounds or skin conditions

Vaccinations and Continuity of Healthcare

Disruption of healthcare services during emergencies may delay vaccinations and medical follow-up, increasing the risk of infectious disease outbreaks. Interruptions in medication supplies or difficulty accessing healthcare facilities may also worsen chronic illnesses and weaken immune defenses.

Routine vaccinations protect children and adults from many serious diseases and should be maintained whenever possible, even during emergencies. Ongoing management of chronic diseases and regular medical follow-up remain essential during crises.

Recommendations include:

- Ensure routine vaccinations are up to date
- Keep vaccination and medical records accessible
- Continue management of chronic illnesses
- Follow prescribed treatment plans

Health-related actions to take:

- Contact healthcare facilities to identify available services
- Keep an adequate supply of essential medications
- Seek medical advice if treatment is interrupted or symptoms worsen
- Consult healthcare providers if exposure to infectious disease is suspected

👉 Continuity of healthcare reduces complications and protects public health.

Disease Prevention in Shelters and Temporary Accommodation Centers

Crowded shelters and temporary accommodation centers may increase the spread of infectious diseases if preventive measures are not followed. These settings may face challenges related to hygiene, ventilation, water supply, and access to healthcare services.

Cooperation between residents, healthcare teams, and emergency responders is essential for maintaining a safe and healthy environment within shelters.

Prevention measures in shelters include:

- Maintaining personal hygiene
- Washing hands regularly
- Avoiding the sharing of personal items
- Reporting symptoms early
- Following instructions from healthcare and emergency teams

Health-related actions to take:

- Isolate sick individuals whenever possible
- Visit healthcare clinics if infectious symptoms appear
- Closely monitor children and older adults
- Use masks when necessary or when recommended by health authorities

 **Cooperation within shelters helps protect everyone's health and safety.**

Protecting Vulnerable Groups

Some groups are more vulnerable to complications from communicable diseases during emergencies due to weakened immunity, chronic illnesses, or difficulty accessing healthcare.

Children, older adults, pregnant women, and people with chronic medical conditions require additional support and continuous health monitoring during emergencies and disasters.

Higher-risk groups include:

- Children
- Older adults
- Pregnant women
- Individuals with chronic diseases
- Immunocompromised individuals

Health-related actions to take:

- Reduce exposure to crowded environments
- Ensure medications and healthcare remain available
- Monitor symptoms closely
- Seek early medical care when symptoms appear

 **Protecting vulnerable groups helps reduce complications and save lives.**

Water Safety During Environmental Emergencies

Environmental emergencies may disrupt water systems or contaminate drinking water sources through flooding, chemical spills, sewage overflow, oil pollution, or infrastructure damage. Unsafe water may carry harmful bacteria, viruses, chemicals, or toxic substances.

Protecting water supplies during emergencies is essential to prevent gastrointestinal illness and other health complications. Families should always follow official water safety advisories and prepare emergency water supplies in advance.

Water safety recommendations:

- Use bottled or approved drinking water if advised
- Avoid drinking water with unusual smell, taste, or color
- Store emergency clean water supplies
- Prevent children from playing in contaminated floodwater
- Follow boil-water notices when issued

Pharmacists can also help explain:

- Medication instructions
- Safe storage
- Side effects
- Drug interactions

 **Safe water practices help reduce infection and illness during emergencies.**

Mental Health and Communicable Diseases

Fear of infection, isolation, uncertainty, and sudden lifestyle changes during emergencies may increase stress, anxiety, and emotional distress. Rumors and misinformation may also negatively affect feelings of safety and emotional well-being.

Mental health is an important part of overall health, as prolonged stress may affect sleep, concentration, relationships, and physical health.

Emotional support recommendations:

- Obtain information from trusted sources
- Avoid spreading rumors or misinformation
- Maintain daily routines whenever possible
- Stay connected with family and community
- Seek psychological support when needed

Health-related actions to take:

- Monitor children and individuals experiencing stress or anxiety
- Encourage supportive family communication
- Seek professional mental health support if distress becomes severe or persistent

👉 Mental health is an essential component of emergency health response and recovery.

Thanks



@EHSUAE
www.ehs.gov.ae